



- (72) DeBonte, Lorin R., US  
(72) Hitz, William D., US  
(71) CARGILL, INCORPORATED, US  
(51) Int.Cl.<sup>6</sup> A01H 5/10, C11B 1/10, C12N 15/53  
(54) **HUILE DE COLZA POSSEDANT UNE TENEUR PLUS ELEVEE  
EN ACIDE OLEIQUE ET EN ACIDE LINOLENIQUE**  
(54) **CANOLA OIL HAVING INCREASED OLEIC ACID AND  
DECREASED LINOLENIC ACID CONTENT**

(57) TRANSLATION NOT AVAILABLE AT THIS  
TIME

(57) An endogenous oil extracted from Brassica seeds is disclosed that contains, after crushing and extraction, greater than 86% oleic acid and less than 2.5% .alpha.-linolenic acid. The oil also contains less than 7% linoleic acid. The Brassica seeds are produced by plants that contain seed-specific inhibition of microsomal oleate desaturase and microsomal linoleate desaturase gene expression. Such inhibition can be created by cosuppression or antisense technology. Such an oil has a very high oxidative stability in the absence of added antioxidants.

2180386

CANOLA OIL HAVING INCREASED OLEIC ACID AND  
DECREASED LINOLENIC ACID CONTENT

ABSTRACT

5 An endogenous oil extracted from *Brassica* seeds is  
disclosed that contains, after crushing and extraction,  
greater than 86% oleic acid and less than 2.5%  $\alpha$ -linolenic  
acid. The oil also contains less than 7% linoleic acid.  
The *Brassica* seeds are produced by plants that contain seed-  
specific inhibition of microsomal oleate desaturase and  
10 microsomal linoleate desaturase gene expression. Such  
inhibition can be created by cosuppression or antisense  
technology. Such an oil has a very high oxidative stability  
in the absence of added antioxidants.

4578.M11

2180386

PATENT  
ATTORNEY DOCKET NO: 87148/042CA1

CANOLA OIL HAVING INCREASED OLEIC ACID AND  
DECREASED LINOLENIC ACID CONTENT

5

Technical Field

This invention relates to a *Brassica* canola oil having an elevated oleic acid content and a decreased linolenic acid profile in the seed oil. The invention also relates to methods by which such an oil may be produced.

10

Background of the Invention

Diets high in saturated fats increase low density lipoproteins (LDL) which mediate the deposition of cholesterol on blood vessels. High plasma levels of serum cholesterol are closely correlated with atherosclerosis and coronary heart disease (Conner et al., *Coronary Heart Disease: Prevention, Complications, and Treatment*, pp. 43-64, 1985). By producing oilseed *Brassica* varieties with reduced levels of individual and total saturated fats in the seed oil, oil-based food products which contain less saturated fats can be produced. Such products will benefit public health by reducing the incidence of atherosclerosis and coronary heart disease.

The dietary effects of monounsaturated fats have also been shown to have dramatic effects on health. Oleic acid, the only monounsaturated fat in most edible vegetable oils, lowers LDL as effectively as linoleic acid, but does not affect high density lipoproteins (HDL) levels (Mattson, F.H., *J. Am. Diet. Assoc.*, 89:387-391, 1989; Mensink et al., *New England J. Med.*, 321:436-441, 1989). Oleic acid is at least as effective in lowering plasma cholesterol as a diet low in fat and high in carbohydrates (Grundy, S.M., *New England J. Med.*, 314:745-748, 1986; Mensink et al., *New England J. Med.*, 321:436-441, 1989). In fact, a high oleic acid diet is preferable to low fat, high carbohydrate diets

2180386

for diabetics (Garg et al., New England J. Med., 319:829-834, 1988). Diets high in monounsaturated fats are also correlated with reduced systolic blood pressure (Williams et al., J. Am. Med. Assoc., 257:3251-3256, 1987).

- 5 Epidemiological studies have demonstrated that the "Mediterranean" diet, which is high in fat and monounsaturates, is not associated with coronary heart disease.

- 10 Intensive breeding has produced *Brassica* plants whose seed oil contains less than 2% erucic acid. The same varieties have also been bred so that the defatted meal contains less than 30  $\mu$ mol glucosinolates/gram. *Brassica* seeds, or oils extracted from *Brassica* seeds, that contain less than 2% erucic acid ( $C_{22:1}$ ), and produce a meal with  
15 less than 30  $\mu$ mol glucosinolates/gram are referred to as canola seeds or canola oils. Plant lines producing such seeds are also referred to as canola lines or varieties.

- Many breeding studies have been directed to alteration of the fatty acid composition in seeds of  
20 *Brassica* varieties. For example, Pleines and Freidt, Fat Sci. Technol., 90(5), 167-171 (1988) describe plant lines with reduced  $C_{18:3}$  levels (2.5-5.8%) combined with high oleic content (73-79%). Roy and Tarr, Z. Pflanzenzuchtg, 95(3), 201-209 (1985) teaches transfer of genes through an  
25 interspecific cross from *Brassica juncea* into *Brassica napus* resulting in a reconstituted line combining high linoleic with low linolenic acid content. Roy and Tarr, Plant Breeding, 98, 89-96 (1987) discuss prospects for development of *B. napus* L. having improved linolenic and linolenic acid  
30 content. Canvin, Can. J. Botany, 43, 63-69 (1965) discusses the effect of temperature on the fatty acid composition of oils from several seed crops including rapeseed.

2180386

Mutations can be induced with extremely high doses of radiation and/or chemical mutagens (Gaul, H. Radiation Botany (1964) 4:155-232). High dose levels which exceed LD50, and typically reach LD90, led to maximum achievable mutation rates. In mutation breeding of *Brassica* varieties, high levels of chemical mutagens alone or combined with radiation have induced a limited number of fatty acid mutations (Rakow, G.Z. Pflanzenzuchtg (1973) 69:62-82).

Rakow and McGregor, J. Amer. Oil Chem. Soc., 50, 400-403 (Oct. 1973) discuss problems associated with selecting mutants affecting seed linoleic and linolenic acid levels. The low  $\alpha$ -linolenic acid mutation derived from the Rakow mutation breeding program did not have direct commercial application because of low seed yield. The first commercial cultivar using the low  $\alpha$ -linolenic acid mutation derived in 1973 was released in 1988 as the variety Stellar (Scarth, R. et al., Can. J. Plant Sci. (1988) 68:509-511). The  $\alpha$ -linolenic acid content of Stellar seeds was greater than 3% and the linoleic acid content was about 28%.

Chemical and/or radiation mutagenesis has been used in an attempt to develop an endogenous canola oil having an oleic acid content of greater than 79% and an  $\alpha$ -linolenic acid content of less than 5%. Wong, et al., EP 0 323 753 B1. However, the lowest  $\alpha$ -linolenic acid level achieved was about 2.7%. PCT publication WO 91/05910 discloses mutagenesis of a starting *Brassica napus* line in order to increase the oleic acid content in the seed oil. However, the oleic acid content in canola oil extracted from seeds of such mutant lines did not exceed 80%.

The quality of canola oil and its suitability for different end uses is in large measure determined by the relative proportion of the various fatty acids present in the seed triacylglycerides. As an example, the oxidative

2180386

stability of canola oil, especially at high temperatures, decreases as the proportion of tri-unsaturated acids increases. Oxidative stability decreases to a lesser extent as the proportion of di-unsaturated acids increases.

- 5 However, it has not been possible to alter the fatty acid composition in *Brassica* seeds beyond certain limits. Thus, an endogenous canola oil having altered fatty acid compositions in seeds is not available for certain specialty uses. Instead, such specialty oils typically are prepared  
10 from canola oil by further processing, such as hydrogenation and/or fractionation.

#### Summary of the Invention

- An endogenous oil obtained from *Brassica* seeds is disclosed. The oil has an oleic acid content of greater  
15 than about 80%, an  $\alpha$ -linolenic acid content of less than about 2.5% and an erucic acid content of less than about 2%, which contents are determined after hydrolysis of the oil. Preferably the oleic acid content is from about 84% to about 88% and the  $\alpha$ -linolenic acid content is from about 1% to  
20 about 2%. The oil may further have a linoleic acid content of from about 1% to about 10%, also determined after hydrolysis of the oil. The oil can be obtained from *Brassica napus* seeds, for example.

- Also disclosed herein is a *Brassica* plant containing  
25 at least one recombinant nucleic construct. The construct(s) comprise a first seed-specific regulatory sequence fragment operably linked to a wild-type microsomal delta-12 fatty acid desaturase coding sequence fragment and a second seed-specific regulatory sequence fragment operably  
30 linked to a wild-type microsomal delta-15 fatty acid desaturase coding sequence fragment. Such a plant produces seeds that yield an oil having an oleic acid content of

2180386

about 86% or greater and an erucic acid content of less than about 2%, which are determined after hydrolysis of the oil.

In some embodiments, the plant contains first and second recombinant nucleic acid constructs, the first construct

5 comprising the delta-12 desaturase coding sequence fragment and the second recombinant nucleic acid construct comprising the delta-15 desaturase coding sequence fragment. The delta-12 or delta-15 desaturase coding sequence fragments may comprise either a partial or a full-length *Brassica*  
10 delta-12 or delta-15 desaturase coding sequence.

Another *Brassica* plant containing at least one recombinant nucleic acid construct is disclosed herein. The construct(s) comprises a first seed-specific regulatory

sequence fragment operably linked to a wild-type microsomal  
15 delta-12 fatty acid desaturase coding sequence fragment and a second seed-specific regulatory sequence fragment operably linked to a wild-type microsomal delta-15 fatty acid

desaturase coding sequence fragment. The plant produces seeds yielding an oil having an oleic acid content of 80% or  
20 greater, an  $\alpha$ -linolenic acid content of about 2.5% or less and an erucic acid content of less than about 2%, which

contents are determined after hydrolysis of the oil. Such a plant may have first and second regulatory sequence fragments linked in sense orientation to the delta-12 and  
25 delta-15 desaturase coding sequence fragments, respectively.

Alternatively the first and second regulatory sequence fragments may be linked in antisense orientation to the corresponding coding sequence fragments. The delta-12 or delta-15 desaturase coding sequence fragments may comprise a

30 partial or a full-length *Brassica* delta-12 or delta-15 desaturase coding sequence. The plant may produce seeds yielding an oil having an oleic acid content of 80% or greater, an  $\alpha$ -linolenic acid content of about 2.5% or less

2180386

and an erucic acid content of less than about 2%, which contents are determined after hydrolysis of the oil.

A method of producing an endogenous oil from Brassica seeds is disclosed herein. The method comprises the steps of: creating at least one Brassica plant having a seed-specific reduction in microsomal delta-12 fatty acid desaturase gene expression and a seed-specific reduction in microsomal delta-15 fatty acid desaturase gene expression; crushing seeds produced from the plant; and extracting the oil from the seeds. The oil has an oleic acid content of about 86% or greater and an erucic acid content of less than about 2%, determined after hydrolysis of the oil. The seed-specific reduction in delta-12 or delta-15 desaturase expression may be created by cosuppression or antisense.

Another method of producing an endogenous oil from Brassica seeds is disclosed herein. The method comprises the steps of: creating at least one Brassica plant having a seed-specific reduction in microsomal delta-12 fatty acid desaturase gene expression and a seed-specific reduction in microsomal delta-15 fatty acid desaturase gene expression; crushing seeds produced from the plant; and extracting the oil from the seeds. The oil has an oleic acid content of about 80% or greater, an  $\alpha$ -linolenic acid content of 2.5% or less and an erucic acid content of less than about 2%, determined after hydrolysis of the oil. The seed-specific reduction in delta-12 or delta-15 desaturase expression may be created by cosuppression or by antisense.

#### Description of the Preferred Embodiments

The term "fatty acid desaturase" refers to an enzyme which catalyzes the breakage of a carbon-hydrogen bond and the introduction of a carbon-carbon double bond into a fatty acid molecule. The fatty acid may be free or esterified to



2180386

another molecule including, but not limited to, acyl-carrier protein, coenzyme A, sterols and the glycerol moiety of glycerolipids. The term "glycerolipid desaturases" refers to a subset of the fatty acid desaturases that act on fatty acyl moieties esterified to a glycerol backbone. "Delta-12 desaturase" refers to a fatty acid desaturase that catalyzes the formation of a double bond between carbon positions 6 and 7 (numbered from the methyl end), (i.e., those that correspond to carbon positions 12 and 13 (numbered from the carbonyl carbon) of an 18 carbon-long fatty acyl chain. "Delta-15 desaturase" refers to a fatty acid desaturase that catalyzes the formation of a double bond between carbon positions 3 and 4 (numbered from the methyl end), (i.e., those that correspond to carbon positions 15 and 16 (numbered from the carbonyl carbon) of an 18 carbon-long fatty acyl chain. "Microsomal desaturase" refers to the cytoplasmic location of the enzyme, while "chloroplast desaturase" and "plastid desaturase" refer to the plastid location of the enzyme. It should be noted that these fatty acid desaturases have never been isolated and characterized as proteins. Accordingly, the terms such as "delta-12 desaturase" and "delta-15 desaturase" are used as a convenience to describe the proteins encoded by nucleic acid fragments that have been isolated based on the phenotypic effects caused by their disruption. They do not imply any catalytic mechanism. For example, delta-12 desaturase refers to the enzyme that catalyzes the formation of a double bond between carbons 12 and 13 of an 18 carbon fatty acid irrespective of whether it "counts" the carbons from the methyl, carboxyl end, or the first double bond.

Microsomal delta-12 fatty acid desaturase (also known as omega-6 fatty acid desaturase, cytoplasmic oleic desaturase or oleate desaturase) is involved in the

enzymatic conversion of oleic acid to linoleic acid. A  
microsomal delta-12 desaturase has been cloned and  
characterized using T-DNA tagging. Okuley, et al., Plant  
Cell 6:147-158 (1994). The nucleotide sequences of higher  
5 plant genes encoding microsomal delta-12 fatty acid  
desaturase are described in Lightner et al., WO94/11516.

Microsomal delta-15 fatty acid desaturase (also  
known as omega-3 fatty acid desaturase, cytoplasmic linoleic  
acid desaturase or linoleate desaturase) is involved in the  
10 enzymatic conversion of linoleic acid to  $\alpha$ -linolenic acid.  
Sequences of higher plant genes encoding microsomal and  
plastid delta-15 fatty acid desaturases are disclosed in  
Yadav, N., et al., Plant Physiol., 103:467-476 (1993), WO  
93/11245 and Arondel, V. et al., Science, 258:1353-1355  
15 (1992).

*Brassica* species have more than one gene for  
endogenous microsomal delta-12 desaturase and more than one  
gene for endogenous microsomal delta-15 desaturase. The  
genes for microsomal delta-12 desaturase are designated *Fad2*  
20 while the genes for microsomal delta-15 desaturase are  
designated *Fad3*. In amphidiploids, each gene is derived  
from one of the ancestral genomes making up the species  
under consideration. The full-length coding sequences for  
the wild-type *Fad2* genes from *Brassica napus* (termed the D  
25 form and the F form) are shown in SEQ ID NO:1 and SEQ ID  
NO:5, respectively. The full-length coding sequence for a  
wild-type *Fad3* gene is disclosed in WO 93/11245.

The inventors have discovered canola oils that have  
novel fatty acid compositions, e.g., very high oleic acid  
30 levels and very low  $\alpha$ -linolenic acid levels. Such oils may  
be obtained by crushing seeds of transgenic *Brassica* plants  
exhibiting a seed-specific reduction in delta-12 desaturase  
and delta-15 desaturase activity; oil of the invention is

2180386

extracted therefrom. Expression of *Fad2* and *Fad3* in seeds is reduced such that the resulting seed oil possesses very high levels of oleic acid and very low levels of  $\alpha$ -linolenic acid. The fatty acid composition of the endogenous seed oil, as determined after hydrolysis of fatty acid esters reflects the novel fatty acid composition of such seeds.

The fatty acid composition of oils disclosed herein is determined by techniques known to the skilled artisan, e.g., hydrolysis of esterified fatty acids (triacylglycerides and the like) in a bulk seed sample followed by gas-liquid chromatography (GLC) analysis of fatty acid methyl esters.

In one embodiment, an oil of the invention has an oleic acid content of about 80% or greater, as well as a surprisingly low  $\alpha$ -linolenic acid content of about 2.5% or less. The oleic acid content is preferably from about 84% to about 89%, more preferably from about 86% to about 89%. The  $\alpha$ -linolenic acid preferably is from about 1% to less than about 2.5%, more preferably from about 1% to about 2%.

The linoleic acid content of an oil of this embodiment typically is less than about 10%, preferably less than about 7%, more preferably from about 2% to about 6%.

Canola seed is crushed by techniques known in the art. The seed typically is tempered by spraying the seed with water to raise the moisture to, for example, 8.5%. The tempered seed is flaked using smooth roller with, for example, a gap setting of 0.23 to 0.27 mm. Heat may be applied to the flakes to deactivate enzymes, facilitate further cell rupturing, coalesce the oil droplets and agglomerate protein particles in order to ease the extraction process.

Typically, oil is removed from the heated canola flakes by a screw press to press out a major fraction of the

2180386

oil from the flakes. The resulting press cake contains some residual oil.

Crude oil produced from the pressing operation typically is passed through a settling tank with a slotted wire drainage top to remove the solids expressed out with the oil in the screw pressing operation. The clarified oil can be passed through a plate and frame filter to remove the remaining fine solid particles.

Canola press cake produced from the screw pressing operation can be extracted with commercial n-Hexane. The canola oil recovered from the extraction process is combined with the clarified oil from the screw pressing operation, resulting in a blended crude oil.

Free fatty acids and gums typically are removed from the crude oil by heating in a batch refining tank to which food grade phosphoric acid has been added. The acid serves to convert the non-hydratable phosphatides to a hydratable form, and to chelate minor metals that are present in the crude oil. The phosphatides and the metal salts are removed from the oil along with the soapstock. The oil-acid mixture is treated with sodium hydroxide solution to neutralize the free fatty acids and the phosphoric acid in the acid-oil mixture. The neutralized free fatty acids, phosphatides, etc. (soapstock) are drained off from the neutralized oil.

A water wash may be done to further reduce the soap content of the oil. The oil may be bleached and deodorized before use, if desired, by techniques known in the art.

A transgenic plant disclosed herein contains at least one recombinant nucleic acid construct. The construct or constructs comprise an oleate desaturase coding sequence fragment and a linoleate desaturase coding sequence fragment, both of which are expressed preferentially in developing seeds. Seed-specific expression of the

2180386

recombinant desaturases results in a seed-specific reduction in native desaturase gene expression. The seed-specific defect in delta-12 and delta-15 desaturase gene expression alters the fatty acid composition in mature seeds produced on the plant, so that the oil obtained from such seeds has the novel fatty acid compositions disclosed herein.

Typically, the oleate and linoleate desaturase sequence fragments are present on separate constructs and are introduced into the non-transgenic parent on separate plasmids. The desaturase fragments may be isolated or derived from, e.g., *Brassica* spp., soybean (*Glycine max*), sunflower and *Arabidopsis*. Preferred host or recipient organisms for introduction of a nucleic acid construct are oil-producing species, such as *Brassica napus*, *B. rapa* and *B. juncea*.

A transgenic plant disclosed herein preferably is homozygous for the transgene containing construct. Such a plant may be used as a parent to develop plant lines or may itself be a member of a plant line, i.e., be one of a group of plants that display little or no genetic variation between individuals for the novel oil composition trait. Such lines may be created by several generations of self-pollination and selection, or vegetative propagation from a single parent using tissue or cell culture techniques. Other means of breeding plant lines from a parent plant are known in the art.

Progeny of a transgenic plant are included within the scope of the invention, provided that such progeny exhibit the novel seed oil characteristics disclosed herein. Progeny of an instant plant include, for example, seeds formed on  $F_1$ ,  $F_2$ ,  $F_3$ , and subsequent generation plants, or seeds formed on  $BC_1$ ,  $BC_2$ ,  $BC_3$ , and subsequent generation plants.

2180386

A seed-specific reduction in *Fad2* and *Fad3* gene expression may be achieved by techniques including, but not limited to, antisense and cosuppression. These phenomena significantly reduce expression of the gene product by the native genes (wild-type or mutated). The reduction in gene expression can be inferred from the decreased level of reaction product and the increased level of substrate in seeds (e.g., decreased 18:2 and increased 18:1), compared to the corresponding levels in plant tissues expressing the native genes.

The preparation of antisense and cosuppression constructs for inhibition of fatty acid desaturases may utilize fragments containing the transcribed sequence for the *Fad2* and *Fad3* fatty acid desaturase genes in canola. These genes have been cloned and sequenced as discussed hereinabove.

Antisense RNA has been used to inhibit plant target genes in a tissue-specific manner. van der Krol et al., *Biotechniques* 6:958-976 (1988). Antisense inhibition has been shown using the entire cDNA sequence as well as a partial cDNA sequence. Sheehy et al., *Proc. Natl. Acad. Sci. USA* 85:8805-8809 (1988); Cannon et al., *Plant Mol. Biol.* 15:39-47 (1990). There is also evidence that 3' non-coding sequence fragment and 5' coding sequence fragments, containing as few as 41 base-pairs of a 1.87 kb cDNA, can play important roles in antisense inhibition. (Ch'ng et al., *Proc. Natl. Acad. Sci. USA* 86:10006-10010 (1989); Cannon et al., *supra*).

The phenomenon of cosuppression has also been used to inhibit plant target genes in a tissue-specific manner. Cosuppression of an endogenous gene using a full-length cDNA sequence as well as a partial cDNA sequence (730 bp of a 1770 bp cDNA) are known. Napoli et al., *The Plant Cell*

2180386

2:279-289 (1990); van der Krol et al., The Plant Cell 2:291-299 (1990); Smith et al., Mol. Gen. Genetics 224:477-481 (1990).

5 Nucleic acid fragments comprising a partial or a full-length delta-12 or delta-15 fatty acid desaturase coding sequence are operably linked to at least one suitable regulatory sequence in antisense orientation (for antisense constructs) or in sense orientation (for cosuppression constructs). Molecular biology techniques for preparing  
10 such chimeric genes are known in the art. The chimeric gene is introduced into a *Brassica* plant and transgenic progeny displaying a fatty acid composition disclosed herein due to antisense or cosuppression are identified. Transgenic plants that produce a seed oil having a fatty acid  
15 composition disclosed herein are selected for use in the invention. Experimental procedures to develop and identify cosuppressed plants involve breeding techniques and fatty acid analytical techniques known in the art.

One may use a partial cDNA sequence for  
20 cosuppression as well as for antisense inhibition. For example, cosuppression of delta-12 desaturase and delta-15 desaturase in *Brassica napus* may be achieved by expressing, in the sense orientation, the entire or partial seed delta-12 desaturase cDNA found in pCF2-165D. See WO 04/11516.

25 Seed-specific expression of native *Fad2* and *Fad3* genes can also be inhibited by non-coding regions of an introduced copy of the gene. See, e.g., Brusslan, J.A. et al. (1993) Plant Cell 5:667-677; Matzke, M.A. et al., Plant Molecular Biology 16:821-830). One skilled in the art can  
30 readily isolate genomic DNA containing sequences that flank desaturase coding sequences and use the non-coding regions for antisense or cosuppression inhibition.

2180386

Regulatory sequences typically do not themselves code for a gene product. Instead, regulatory sequences affect the expression level of the mutant coding sequence. Examples of regulatory sequences are known in the art and include, without limitation, promoters of genes expressed during embryogenesis, e.g., a napin promoter, a phaseolin promoter, a oleosin promoter and a cruciferin promoter. Native regulatory sequences, including the native promoters, of delta-12 and delta-15 fatty acid desaturase genes can be readily isolated by those skilled in the art and used in constructs of the invention. Other examples of suitable regulatory sequences include enhancers or enhancer-like elements, introns and 3' non-coding regions such as poly A sequences. Further examples of suitable regulatory sequences for the proper expression of mutant or wild-type delta-12 or mutant delta-15 coding sequences are known in the art.

In preferred embodiments, regulatory sequences are seed-specific, i.e., the chimeric desaturase gene product is preferentially expressed in developing seeds and expressed at low levels or not at all in the remaining tissues of the plant. Seed-specific regulatory sequences preferably stimulate or induce expression of the recombinant desaturase coding sequence fragment at a time that coincides with or slightly precedes expression of the native desaturase gene. Murphy et al., J. Plant Physiol. 135:63-69 (1989).

Transgenic plants for use in the invention are created by transforming plant cells of *Brassica* species. Such techniques include, without limitation, *Agrobacterium*-mediated transformation, electroporation and particle gun transformation. Illustrative examples of transformation techniques are described in U.S. Patent 5,204,253, (particle gun) and U.S. Patent 5,188,958 (*Agrobacterium*), incorporated



2180386

herein by reference. Transformation methods utilizing the Ti and Ri plasmids of *Agrobacterium* spp. typically use binary type vectors. Walkerpeach, C. et al., in *Plant Molecular Biology Manual*, S. Gelvin and R. Schilperoort, eds., Kluwer Dordrecht, C1:1-19 (1994). If cell or tissue cultures are used as the recipient tissue for transformation, plants can be regenerated from transformed cultures by techniques known to those skilled in the art.

One or more recombinant nucleic acid constructs, suitable for antisense or cosuppression of native *Fad2* and *Fad3* genes are introduced, and at least one transgenic *Brassica* plant is obtained. Seeds produced by the transgenic plant(s) are grown and either selfed or outcrossed to obtain plants homozygous for the recombinant construct. Seeds are analyzed as discussed above in order to identify those homozygotes having native fatty acid desaturase activities inhibited by the mechanisms discussed above. Homozygotes may be entered into a breeding program, e.g., to increase seed, to introgress the novel oil composition trait into other lines or species, or for further selection of other desirable traits (disease resistance, yield and the like).

Fatty acid composition is followed during the breeding program by analysis of a bulked seed sample or of a single half-seed. Half-seed analysis useful because the viability of the embryo is maintained and thus those seeds having a desired fatty acid profile may be advanced to the next generation. However, half-seed analysis is also known to be an inaccurate representation of the genotype of the seed being analyzed. Bulk seed analysis typically yields a more accurate representation of the fatty acid profile in seeds of a given genotype.

Procedures for analysis of fatty acid composition are known in the art. These procedures can be used to

2180386

identify individuals to be retained in a breeding program; the procedures can also be used to determine the product specifications of commercial or pilot plant oils.

The relative content of each fatty acid in canola seeds can be determined either by direct trans-esterification of individual seeds in methanolic  $H_2SO_4$  (2.5%) or by hexane extraction of bulk seed samples followed by trans-esterification of an aliquot in 1% sodium methoxide in methanol. Fatty acid methyl esters can be extracted from the methanolic solutions into hexane after the addition of an equal volume of water.

For example, a seed sample from each transformant in a breeding program is crushed with a mortar and pestle and extracted 4 times with 8 mL hexane at about 50°C. The extracts from each sample are reduced in volume and two aliquots are taken for esterification. Separation of the fatty acid methyl esters can be carried out by gas-liquid chromatography using an Omegawax 320 column (Supelco Inc., 0.32 mm ID X 30M) run isothermally at 220° and cycled to 260° between each injection.

Alternatively, seed samples from a breeding program are ground and extracted in methanol/KOH, extracted with iso-octane, and fatty acids separated by gas chromatography.

A method to produce an oil of the invention comprises the creation of at least one *Brassica* plant having a seed-specific reduction in *Fad2* and *Fad3* gene expression, as discussed above. Seeds produced by such a plant, or its progeny, are crushed and the oil is extracted from the crushed seeds. Such lines produce seeds yielding an oil of the invention, e.g., an oil having from about 80% to about 88% oleic acid, from about 1% to about 2%  $\alpha$ -linolenic acid and less than about 2% erucic acid.

2180386

Alternatively, such a plant can be created by crossing two parent plants, one exhibiting a reduction in *Fad2* gene expression and the other exhibiting a reduction in *Fad3* gene expression. Progeny of the cross are outcrossed or selfed in order to obtain progeny seeds homozygous for both traits.

Transgenic plants having a substantial reduction in *Fad2* and *Fad3* gene expression in seeds have novel fatty acid profiles in oil extracted from such seeds, compared to known canola plants, e.g., the reduction in both desaturase activities results in a novel combination of high oleic and lower  $\alpha$ -linolenic acid in seed oils. By combining seed-specific inhibition of microsomal delta-12 desaturase with seed-specific inhibition of microsomal delta-15 desaturase, one obtains very low levels of seed  $\alpha$ -linolenic acid, without adversely affecting agronomic properties.

It is noteworthy that *Fad2* and *Fad3* cosuppression constructs provide a novel means for producing canola oil having 86% oleic acid or greater. A method of producing a canola oil having greater than 86% oleic acid comprises the creation of a transgenic *Brassica* plant containing at least one recombinant nucleic acid construct, which construct(s) comprises an oleate desaturase coding sequence expressed preferentially in developing seeds and a linoleate desaturase coding sequence expressed preferentially in developing seeds. A proportion of the plants that are homozygous for the transgenes have seed-specific cosuppression of the native linoleate desaturase. Seeds produced by such transgenic cosuppressed plants are crushed and the oil is extracted therefrom. The oil has about 86% or greater oleic acid and less than about 2% erucic acid. The oleic acid content can be as high as 89%.

2180386

Transgenic plants exhibiting cosuppression of *Fad2* and *Fad3* produce seeds having a very high oleic acid content. This result was unexpected because it was not known if one could obtain plants in which inhibition of *Fad2* and *Fad3* via cosuppression was sufficient to achieve an oleic acid level of 86% or greater in seeds. Indeed, it was not known if two cosuppressed genes in fatty acid metabolism could be introduced in canola without the first cosuppression gene interfering with the second cosuppression gene, or without adversely affecting other agronomic traits.

Marker-assisted breeding techniques may be used to identify and follow a desired fatty acid composition during the breeding process. Such markers may include RFLP, RAPD, or PCR markers, for example. Marker-assisted breeding techniques may be used in addition to, or as an alternative to, other sorts of identification techniques. An example of marker-assisted breeding is the use of PCR primers that specifically amplify the junction between a promoter fragment and the coding sequence of a *Fad2* gene.

While the invention is susceptible to various modifications and alternative forms, certain specific embodiments thereof are described in the general methods and examples set forth below. For example the invention may be applied to all *Brassica* species, including *B. rapa*, *B. juncea*, and *B. hirta*, to produce substantially similar results. It should be understood, however, that these examples are not intended to limit the invention to the particular forms disclosed. Instead, the disclosure is to cover all modifications, equivalents and alternatives falling within the scope of the invention.

#### EXAMPLE 1

#### CONSTRUCTS FOR COSUPPRESSION OF DELTA-12 FATTY

2180386

ACID DESATURASE AND DELTA-15 FATTY ACID DESATURASE

The wild-type *Brassica* cDNA coding sequence for the delta-12 desaturase D form was cloned as described in WO 94/11516, which is incorporated herein by reference.

5 Briefly, rapeseed cDNAs encoding cytoplasmic oleate (18:1) desaturase were obtained by screening a cDNA library made from developing rapeseed using a heterologous probe derived from an *Arabidopsis* cDNA fragment encoding the same enzyme. (Okuley et al 1994). The full-length coding sequence of  
10 *Fad2* is found as SEQ ID NO:1. Rapeseed cDNAs encoding the cytoplasmic linoleate (18:2) desaturase (*Fad3*) were obtained as described in WO 93/11245, incorporated herein by reference. See also (Yadav et. al 1993). Seed specific expression of these cDNAs in transgenic rapeseed was driven  
15 by one of four different seed storage protein promoters, napin, oleosin and cruciferin promoters from *B. napus* and a phaseolin promoter from *Phaseolus vulgaris*.

Detailed procedures for manipulation of DNA fragments by restriction endonuclease digestion, size  
20 separation by agarose gel electrophoresis, isolation of DNA fragments from agarose gels, ligation of DNA fragments, modification of cut ends of DNA fragments and transformation of *E. coli* cells with plasmids have been described. Sambrook et al., (Molecular Cloning, A Laboratory Manual,  
25 2nd ed (1989) Cold Spring Harbor Laboratory Press); Ausubel et al., Current Protocols in Molecular Biology (1989) John Wiley & Sons). Plant molecular biology procedures are described in Plant Molecular Biology Manual, Gelvin S. and Schilperoort, R. eds. Kluwer, Dordrecht (1994).

30 The plasmid pZS212 was used to construct binary vectors for these experiments. pZS212 contains a chimeric CaMV35S/NPT gene for use in selecting kanamycin resistant

2180386

transformed plant cells, the left and right border of an *Agrobacterium* Ti plasmid T-DNA, the *E. coli* lacZ  $\alpha$ -complementing segment with unique restriction endonuclease sites for EcoRI, KpnI, BamHI and SalI, the bacterial  
5 replication origin from the *Pseudomonas* plasmid pVS1 and a bacterial Tn5 NPT gene for selection of transformed *Agrobacterium*. See WO 94/11516, p. 100.

The first construct was prepared by inserting a full-length mutant *Brassica* Fad2 D gene coding sequence  
10 fragment in sense orientation between the phaseolin promoter and phaseolin 3' poly A region of plasmid pCW108. The full-length coding sequence of the mutant gene is found in SEQ ID NO:3.

The pCW108 vector contains the bean phaseolin  
15 promoter and 3' untranslated region and was derived from the commercially available pUC18 plasmid (Gibco-BRL) via plasmids AS3 and pCW104. Plasmid AS3 contains 495 base pairs of the *Phaseolus vulgaris* phaseolin promoter starting with 5'-TGGTCTTTTGGT-3' followed by the entire 1175 base  
20 pairs of the 3' untranslated region of the same gene. Sequence descriptions of the 7S seed storage protein promoter are found in Doyle et al., J. Biol. Chem. 261:9228-9238 (1986) and Slightom et al., Proc. Natl. Acad. Sci. USA, 80:1897-1901 (1983). Further sequence description may be  
25 found in WO 91/13993. The fragment was cloned into the Hind III site of pUC18. The additional cloning sites of the pUC18 multiple cloning region (Eco RI, Sph I, Pst I and Sal I) were removed by digesting with Eco RI and Sal I, filling in the ends with Klenow and religating to yield the plasmid  
30 pCW104. A new multiple cloning site was created between the 495bp of the 5' phaseolin and the 1175bp of the 3' phaseolin by inserting a dimer of complementary synthetic oligonucleotides to create the plasmid pCW108. See WO

2180386

94/11516. This plasmid contains unique Nco I, Sma I, Kpn I and Xba I sites directly behind the phaseolin promoter.

The phaseolin promoter:mutantFad2:phaseolin poly A construct in pCW108 was excised and cloned between the 5 SalI/EcoRI sites of pZS212. The resulting plasmid was designated pIMC201.

A second plasmid was constructed by inserting the full-length wild type *Brassica* Fad2 D gene coding sequence into the NotI site of plasmid pIMC401, which contains a 2.2 10 kb napin expression cassette. See, e.g., WO94/11516, page 102. The 5'-napin:Fad2:napin poly A-3' construct was inserted into the SalI site of pZS212 and the resulting 17.2 Kb plasmid was termed pIMC127. Napin promoter sequences are also disclosed in U.S. Patent 5,420,034.

15 A third plasmid, pIMC135, was constructed in a manner similar to that described above for pIMC127. Plasmid pIMC135 contains a 5' cruciferin promoter fragment operably linked in sense orientation to the full-length wild-type *Brassica* Fad2 D gene coding sequence, followed by a 20 cruciferin 3' poly A fragment. The 5'-cruciferin:Fad2 D:cruciferin polyA cassette was inserted into pZS212; the resulting plasmid was termed pIMC135. Suitable cruciferin regulatory sequences are disclosed in Rodin, J. et al., J. Biol. Chem. 265:2720 (1990); Ryan, A. et al., Nucl. Acids 25 Res. 17:3584 (1989) and Simon, A. et al., Plant Mol. Biol. 5:191 (1985). Suitable sequences are also disclosed in the Genbank computer database, e.g., Accession No. M93103.

A fourth plasmid, pIMC133 was constructed in a manner similar to that described above. Plasmid pIMC133 30 contains a 5' oleosin promoter fragment operably linked in sense orientation to the full-length *Brassica* Fad2 D gene coding sequence, followed by a 3' oleosin poly A fragment. See, e.g., WO 93/20216, incorporated herein by reference.

2180386

A napin-Fad3 construct was made by first isolating a delta-15 desaturase coding sequence fragment from pBNS73-f2. The fragment contained the full-length coding sequence of the desaturase, disclosed as SEQ ID NO: 6 in WO 93/11245, incorporated herein by reference. The 1.2 kb fragment was fitted with linkers and ligated into pIMC401. The 5'napin:Fad3:3'napin cassette was inserted into the Sal I site of pZS212; the resulting plasmid was designated pIMC110.

## EXAMPLE 2

### CREATION OF TRANSGENIC COSUPPRESSED PLANTS

The plasmids pIMC201, pIMC127, pIMC135, pIMC133 and pIMC110 were introduced into *Agrobacterium* strain LBA4404/pAL4404 by a freeze-thaw method. The plasmids were introduced into *Brassica napus* cultivar Westar by the method of *Agrobacterium*-mediated transformation as described in WO94/11516, incorporated herein by reference. Transgenic progeny plants containing pIMC201 were designated as the WS201 series. Plants transformed with pIMC127 were designated as the WS687 series. Plants transformed with pIMC135 were designated as the WS691 series. Plants transformed with pIMC133 were designated as the WS692 series. Plants transformed with pIMC110 were designated as the WS663 series.

Unless indicated otherwise, fatty acid percentages described herein are percent by weight of the oil in the indicated seeds as determined after extraction and hydrolysis.

From about 50 to 350 transformed plants (T1 generation) were produced for each cDNA and promoter combination. T1 plants were selfed to obtain T2 seed. T2 samples in which cosuppression events occurred were



2180386

identified from the fatty acid profile and from the presence of the transgene by molecular analysis. The transformed plants were screened for phenotype by analysis of the relative fatty acid contents of bulk seed from the first transformed generation by GC separation of fatty acid methyl esters.

T2 seed was sown in 4-inch pots containing Pro-Mix soil. The plants, along with Westar controls, were grown at  $25\pm 3^{\circ}\text{C}/18\pm 3^{\circ}\text{C}$ , 14/10 hr day/night conditions in the greenhouse. At flowering, the terminal raceme was self-pollinated by bagging. At maturity, seed was individually harvested from each plant, labelled, and stored to ensure that the source of the seed was known.

Fatty acid profiles were determined as described in WO 91/05910. For chemical analysis, 10-seed bulk samples were hand ground with a glass rod in a 15-mL polypropylene tube and extracted in 1.2 mL 0.25 N KOH in 1:1 ether/methanol. The sample was vortexed for 30 sec. and heated for 60 sec. in a  $60^{\circ}\text{C}$  water bath. Four mL of saturated NaCl and 2.4 mL of iso-octane were added, and the mixture was vortexed again. After phase separation, 600  $\mu\text{L}$  of the upper organic phase were pipetted into individual vials and stored under nitrogen at  $-5^{\circ}\text{C}$ . One  $\mu\text{L}$  samples were injected into a Supelco SP-2330 fused silica capillary column (0.25 mm ID, 30 M length, 0.20  $\mu\text{m}$  df).

The gas chromatograph was set at  $180^{\circ}\text{C}$  for 5.5 minutes, then programmed for a  $2^{\circ}\text{C}/\text{minute}$  increase to  $212^{\circ}\text{C}$ , and held at this temperature for 1.5 minutes. Total run time was 23 minutes. Chromatography settings were: Column head pressure - 15 psi, Column flow (He) - 0.7 mL/min., Auxiliary and Column flow - 33 mL/min., Hydrogen flow - 33 mL/min., Air flow - 400 mL/min., Injector temperature -  $250^{\circ}\text{C}$ , Detector temperature -  $300^{\circ}\text{C}$ , Split vent - 1/15.

2180386

Table 1 shows the content of the seven major fatty acids in mature seeds from transgenic cosuppressed plants homozygous for the napin:Fad3 construct or the napin:Fad2 construct (T4 or later generation). Over expression phenotypes and cosuppression phenotypes were observed for both chimeric genes (oleate desaturase and linoleate desaturase); data for plants exhibiting the cosuppression phenotype are shown in the Table.

As shown in Table 1, the homozygous Fad2-cosuppressed seed had a  $\alpha$ -linolenic acid content of about 2.9%, which was less than half that of the Westar control; the oleic acid content increased to about 84.1%. The homozygous Fad3-cosuppressed seed had an  $\alpha$ -linolenic acid of about 1.2%; the oleic acid and linoleic acid contents in Fad3-cosuppressed plants increased slightly compared to Westar. The results demonstrate that inhibiting gene expression of either enzyme by cosuppression resulted in a change in fatty acid composition of the seed oil.

TABLE 1

Fatty Acid Profiles in Oil From Cosuppression Canola Seed

TRANSGENE CONSTRUCTION	FATTY ACID (% OF TOTAL FATTY ACIDS)								
	16:0	18:0	18:1	18:2	18:3	20:0	20:1	22:0	24:0
non-transformed Westar	3.9	1.6	67.0	19.0	7.5	0.6	0.8	0.6	0.1
napin:Fad2 (co-suppression)	4.3	1.4	84.1	5.2	2.9	0.6	0.9	0.5	0.2
napin:Fad3 (co-suppression)	3.8	1.5	68.5	22.1	1.2	0.6	1.1	0.4	0.1

2180386

TABLE 2

Fatty Acid Profiles in Oil From Cosuppression Canola Seeds

Line #	Construct (promoter/coding sequence)	Fatty Acid Composition				
		16:0	18:0	18:1	18:2	18:3
5	663-40 napin/ <i>Fad3</i>	3.9	1.4	71.2	20.1	1.2
	687-193 napin/ <i>Fad2</i>	4.0	1.5	82.8	5.9	3.7
	691-215 cruciferin/ <i>Fad2</i>	3.3	1.3	86.5	3.0	3.7
	692-090-3 oleosin/ <i>Fad2</i>	3.4	1.3	86.5	2.6	3.9
	692-105-11 oleosin/ <i>Fad2</i>	3.4	1.3	86.2	2.7	4.2
10	201-389 A23 phaseolin/M <i>Fad2</i>	4.2	2.7	84.6	4.7	3.7

TABLE 3

Range of Fatty Acid Profiles for *Fad2* and *Fad3*  
Cosuppression Lines Tested in the Field

Line No.	Vector	Min/Max	Fatty Acid Composition				
			C16:0	C18:0	C18:1	C18:2	C18:3
15	663-40 pIMC110	Min	3.5	2.3	73.5	16.3	0.8
		Max	4.7	2.2	64.0	24.2	1.5
	687-193 pIMC127	Min	3.4	3.1	83.3	3.8	2.3
		Max	3.4	2.1	85.5	3.2	2.5
20	692-105 pIMC133	Min	3.7	2.7	84.6	2.8	2.4
		Max	3.3	2.3	86.3	2.1	2.7
	691-215 pIMC135	Min	3.2	2.4	84.6	3.0	2.5
		Max	3.0	2.0	86.3	2.6	2.5

Table 2 shows the fatty acid profile in T4 or later homozygous seeds produced by six individual plants having various promoter-desaturase gene combinations. The seeds were obtained from greenhouse-grown plants. The results indicate that the oleic acid content ranged from about 82.8% to about 86.5% among the lines carrying the *Fad2* constructs. The phaseolin:mutated *Fad2* construct was as successful as the wild-type *Fad2* constructs in achieving seed-specific *Fad2* cosuppression.

The napin:*Fad3* cosuppressed plant line had an unusually low  $\alpha$ -linolenic acid content of 1.2%. However, the oleic acid content was only 71.2% and the linoleic acid

2180386

content was similar to that of the non-transformed control Westar in Table 1.

5       Homozygous seeds from four of the lines in Table 2 were planted in a field nursery in Colorado and self-pollinated. Seed samples from several plants of each line were collected and separately analyzed for fatty acid composition. The results for the 663-40 plant having the minimum and the 663-40 plant having the maximum linolenic acid content observed in the field are shown in Table 3.

10      The results for the 687-193, 692-105 and 691-215 plants having the minimum and maximum oleic acid content in the field are also shown in Table 3.

15       The results in Table 3 demonstrate that the fatty acid profile in field-grown seeds of cosuppressed transgenic plants was similar to that in the greenhouse-grown seeds (Table 2), indicating that the cosuppression trait confers a stable fatty acid composition on the oil. The results also indicate that an oil having the combination of an oleic acid content of 86% or greater and an  $\alpha$ -linolenic acid content of

20      2.5% or less could not be obtained from plants cosuppressed for either *Fad2* or *Fad3* alone.

### EXAMPLE 3

#### OIL CONTENT IN SEEDS OF PLANTS EXHIBITING *Fad2* and *Fad3* COSUPPRESSION

25       Crosses were made between the napin:*Fad3* cosuppressed line 663-40 and three *Fad2* cosuppressed lines, 691-215, 692-090-3 and 692-105-11. F1 plants were selfed for 2 generations in the greenhouse to obtain F3 generation seed that was homozygous for both recombinant constructs.

2180386

TABLE 4

Fatty Acid Profile in F3 Seeds of  
Lines Exhibiting Fad2 and Fad3 Cosuppression

Line #	Construct	16:0	18:0	18:1	18:2	18:3
5	663-40 napin/Fad3	3.9	1.4	71.2	20.1	1.2
	691-215 cruciferin/Fad2	3.9	1.3	86.5	3.0	3.7
	663-40X691-215 napin/Fad3 & cruciferin/Fad2	3.2	1.4	86.2	5.2	1.5
		16:0	18:0	18:1	18:2	18:3
10	663-40 napin/Fad3	3.9	1.4	71.2	20.1	1.2
	692-090-3 oleosin/Fad2	3.4	1.3	86.5	2.6	3.9
	663-40X692-090-3 napin/Fad3 & oleosin/Fad2	3.4	1.5	85.5	5.0	1.7
		16:0	18:0	18:1	18:2	18:3
15	663-40 napin/Fad3	3.9	1.4	71.2	20.1	1.2
	692-105-11 oleosin/Fad2	3.4	1.3	86.2	2.7	4.2
	663-40X692-105-11 napin/Fad3 & oleosin/Fad2	3.4	1.4	86.8	4.6	1.4

The seed fatty acid profiles of the parent lines and a representative F3 cosuppressed line are shown in Table 4. Plants expressing both cosuppression constructs exhibited an oleic acid level of about 86% or greater. Moreover, this high level of oleic acid was present in combination with an unusually low level of  $\alpha$ -linolenic acid, less than 2.0%. However, the linoleic acid content in the F3 seeds increased from about 2.6-3.0% to about 4.6-5.2%.

These results demonstrate that a canola oil can be extracted from rapeseeds that contains greater than 80% oleic acid and less than 2.5%  $\alpha$ -linolenic acid. Results similar to those obtained using cosuppression constructs are achieved when antisense constructs are used.

The canola oil extracted from Fad2 and Fad3 cosuppressed F3 seed, or progeny thereof, is found to have superior oxidative stability compared to the oil extracted from Westar seed. The improved oxidative stability of such an oil is measured after refining, bleaching and deodorizing, using the Accelerated Oxygen Method (AOM),

2180386

American Oil Chemists' Society Official Method Cd 12-57 for fat stability, Active Oxygen Method (revised 1989). The improved oxidative stability is also demonstrated when using the Oxidative Stability Index method. The improved oxidative stability is measured in the absence of added antioxidants.

#### EXAMPLE 4

##### OIL CONTENT IN SEEDS OF PLANTS HAVING *Fad3* COSUPPRESSION AND CHEMICALLY-INDUCED *Fad2* MUTATIONS

Q4275 is a doubly mutagenized *B. napus* line having defects in the *Fad2* gene. Q4275 was derived by chemical mutagenesis of *B. napus* line IMC129, which carries a mutation in the *Fad2* D gene; the coding sequence of the mutated gene is shown in SEQ ID NO:3. Line IMC129 was itself derived by chemical mutagenesis of the cultivar Westar, as disclosed in WO 91/05910. Genetic segregation analysis of crosses between Q4275 and other fatty acid mutant lines indicated that Q4275 carried a mutation in the *B. napus* *Fad2* F gene in addition to the IMC129 *Fad2* D gene mutation. Q4275 thus carries chemically induced mutations in both *Fad2* genes.

A cross was made between Q4275 and the napin:*Fad3* cosuppressed line 663-40. F1 plants were selfed in the greenhouse and F2 plants that were homozygous for the recombinant construct and the *Fad2* D and *Fad2* F mutated genes were identified by fatty acid profile analysis of the F3 generation seed. After selfing to homozygosity, the fatty acid profiles in seeds of a representative homozygous plant was analyzed and compared to the profile of the parent plants, as shown in Table 5.

The results show that an oil having greater than 87% oleic acid and less than 1.5%  $\alpha$ -linolenic acid can be

2180386

obtained from a transgenic *Brassica* plant containing a seed-specific reduction in *Fad3* gene expression as well as chemically-induced mutations in *Fad2* genes.

TABLE 5

5 Fatty Acid Profile of *Fad3* Cosuppression, *Fad2* Mutated Seeds

	16:0	18:0	18:1	18:2	18:3
663-40	3.9	1.4	71.2	20.1	1.2
Q4275	3.3	1.5	86.7	2.2	3.1
Q4275 X 663-40	3.2	1.6	87.6	4.2	1.3

10

TABLE 6

Range of Fatty Acid Profiles for *Fad3* Cosuppression,  
*Fad2* Mutated Lines Tested in the Field

		16:0	18:0	18:1	18:2	18:3
15	663-40	Min	3.5	2.3	73.5	16.3
		Max	4.7	2.2	64.0	24.2
	Q4275	Min	3.2	3.3	85.0	1.8
		Max	3.0	2.3	86.6	1.7
	Q4275 X 663-40	Min	3.2	2.0	85.1	5.3
		Max	3.2	2.9	84.0	6.0

20 Additional seed from the homozygous plant described above was planted in the field and self-pollinated. Mature seeds from several progeny plants were separately analyzed for their fatty acid profile. The fatty acid profile for the progeny plant having the minimum linolenic acid content and the plant having the maximum linolenic acid content are shown in Table 6. The results show that the homozygous plant having *Fad2* mutations and *Fad3* cosuppression had a fatty acid profile in the field that was similar to that of

25

2180386

the greenhouse-grown seed (Table 5), indicating that the *Fad3* cosuppression trait and the chemically-induced *Fad2* mutants conferred a stable fatty acid composition on seeds of this plant. Thus, an oil of the invention can be  
5 obtained from either field-grown seeds or greenhouse-grown seeds.

Because of the decreased  $\alpha$ -linolenic acid content and increased oleic acid content, an oil of the invention is useful in food and industrial applications. Oils which are  
10 low in  $\alpha$ -linolenic acid have increased oxidative stability. The rate of oxidation of lipid fatty acids increases with higher levels of linolenic acid leading to off-flavors and off-odors in foods. The present invention provides novel canola oils that are low in  $\alpha$ -linolenic acid.

15 To the extent not already indicated, it will be understood by those of ordinary skill in the art that any one of the various specific embodiments herein described and illustrated may be further modified to incorporate features shown in other of the specific embodiments.

20 The foregoing detailed description has been provided for a better understanding of the invention only and no unnecessary limitation should be understood therefrom as some modifications will be apparent to those skilled in the art without deviating from the spirit and scope of the  
25 appended claims.



2180386

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: DeBonte, L. et al.
- (ii) TITLE OF INVENTION: CANOLA OIL HAVING INCREASED OLEIC ACID AND DECREASED LINOLENIC ACID CONTENT
- (iii) NUMBER OF SEQUENCES: 6
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Fish & Richardson, P.C., P.A.
  - (B) STREET: 60 South Sixth Street, Suite 3300
  - (C) CITY: Minneapolis
  - (D) STATE: MN
  - (E) COUNTRY: USA
  - (F) ZIP: 55402
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE: 03-JUL-1996
  - (C) CLASSIFICATION:
- (vii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Lundquist, Ronald C.
  - (B) REGISTRATION NUMBER: 37,875
  - (C) REFERENCE/DOCKET NUMBER: 07148/042001
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: 612/335-5070
  - (B) TELEFAX: 612/288-9696

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1155 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Brassica napus
- (ix) FEATURE:
  - (D) OTHER INFORMATION: Wild type D form.

2180386

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATG GGT GCA GGT GGA AGA ATG CAA GTG TCT CCT CCC TCC AAG AAG TCT	48
Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser	
1 5 10 15	
GAA ACC GAC ACC ATC AAG CGC GTA CCC TGC GAG ACA CCG CCC TTC ACT	96
Glu Thr Asp Thr Ile Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr	
20 25 30	
GTC GGA GAA CTC AAG AAA GCA ATC CCA CCG CAC TGT TTC AAA CGC TCG	144
Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser	
35 40 45	
ATC CCT CGC TCT TTC TCC TAC CTC ATC TGG GAC ATC ATC ATA GCC TCC	192
Ile Pro Arg Ser Phe Ser Tyr Leu Ile Trp Asp Ile Ile Ile Ala Ser	
50 55 60	
TGC TTC TAC TAC NTC GCC ACC ACT TAC TTC CCT CTC CTC CCT CAC CCT	240
Cys Phe Tyr Tyr Xaa Ala Thr Thr Tyr Phe Pro Leu Leu Pro His Pro	
65 70 75 80	
CTC TCC TAC TTC GCC TGG CCT CTC TAC TGG GCC TGC CAA GGG TGC GTC	288
Leu Ser Tyr Phe Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val	
85 90 95	
CTA ACC GGC GTC TGG GTC ATA GCC CAC GAA TGC GGC CAC CAC GCC TTC	336
Leu Thr Gly Val Trp Val Ile Ala His Glu Cys Gly His His Ala Phe	
100 105 110	
AGC GAC TAC CAG TGG CTT GAC GAC ACC GTC GGT CTC ATC TTC CAC TCC	384
Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser	
115 120 125	
TTC CTC CTC GTC CCT TAC TTC TCC TGG AAG TAC AGT CAT CGC AGC CAC	432
Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Ser His	
130 135 140	
CAT TCC AAC ACT GGC TCC CTC GAG AGA GAC GAA GTG TTT GTC CCC AAG	480
His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys	
145 150 155 160	
AAG AAG TCA GAC ATC AAG TGG TAC GGC AAG TAC CTC AAC AAC CCT TTG	528
Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu	
165 170 175	
GGA CGC ACC GTG ATG TTA ACG GTT CAG TTC ACT CTC GGC TGG CCG TTG	576
Gly Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu	
180 185 190	
TAC TTA GCC TTC AAC GTC TCG GGA AGA CCT TAC GAC GGC GGC TTC CGT	624
Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Arg	
195 200 205	
TGC CAT TTC CAC CCC AAC GCT CCC ATC TAC AAC GAC CGC GAG CGT CTC	672
Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Arg Glu Arg Leu	
210 215 220	
CAG ATA TAC ATC TCC GAC GCT GGC ATC CTC GCC GTC TGC TAC GGT CTC	720
Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu	
225 230 235 240	

2180386

TTC CGT TAC GCC GCC GGC CAG GGA GTG GCC TCG ATG GTC TGC TTC TAC Phe Arg Tyr Ala Ala Gly Gln Gly Val Ala Ser Met Val Cys Phe Tyr 245 250 255	768
GGA GTC CCG CTT CTG ATT GTC AAT GGT TTC CTC GTG TTG ATC ACT TAC Gly Val Pro Leu Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr 260 265 270	816
TTG CAG CAC ACG CAT CCT TCC CTG CCT CAC TAC GAT TCG TCC GAG TGG Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp 275 280 285	864
GAT TGG TTC AGG GGA GCT TTG GCT ACC GTT GAC AGA GAC TAC GGA ATC Asp Trp Phe Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile 290 295 300	912
TTG AAC AAG GTC TTC CAC AAT ATT ACC GAC ACG CAC GTG GCC CAT CAT Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His 305 310 315 320	960
CCG TTC TCC ACG ATG CCG CAT TAT CAC GCG ATG GAA GCT ACC AAG GCG Pro Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala 325 330 335	1008
ATA AAG CCG ATA CTG GGA GAG TAT TAT CAG TTC GAT GGG ACG CCG GTG Ile Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Phe Asp Gly Thr Pro Val 340 345 350	1056
GTT AAG GCG ATG TGG AGG GAG GCG AAG GAG TGT ATC TAT GTG GAA CCG Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro 355 360 365	1104
GAC AGG CAA GGT GAG AAG AAA GGT GTG TTC TGG TAC AAC AAT AAG TTA T Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu 370 375 380	1153
GA	1155

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 384 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser 1 5 10 15
Glu Thr Asp Thr Ile Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr 20 25 30
Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser 35 40 45
Ile Pro Arg Ser Phe Ser Tyr Leu Ile Trp Asp Ile Ile Ile Ala Ser 50 55 60

2180386

Cys Phe Tyr Tyr Xaa Ala Thr Thr Tyr Phe Pro Leu Leu Pro His Pro  
 65 70 75 80  
 Leu Ser Tyr Phe Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val  
 85 90 95  
 Leu Thr Gly Val Trp Val Ile Ala His Glu Cys Gly His His Ala Phe  
 100 105 110  
 Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser  
 115 120 125  
 Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Ser His  
 130 135 140  
 His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys  
 145 150 155 160  
 Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu  
 165 170 175  
 Gly Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu  
 180 185 190  
 Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Arg  
 195 200 205  
 Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu  
 210 215 220  
 Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu  
 225 230 235 240  
 Phe Arg Tyr Ala Ala Gly Gln Gly Val Ala Ser Met Val Cys Phe Tyr  
 245 250 255  
 Gly Val Pro Leu Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr  
 260 265 270  
 Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp  
 275 280 285  
 Asp Trp Phe Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile  
 290 295 300  
 Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His  
 305 310 315 320  
 Pro Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala  
 325 330 335  
 Ile Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Phe Asp Gly Thr Pro Val  
 340 345 350  
 Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro  
 355 360 365  
 Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu  
 370 375 380

2180386

## (2) INFORMATION FOR SEQ ID NO:3:

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1155 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Brassica napus*

(vii) IMMEDIATE SOURCE:

(B) CLONE: IMC129

(ix) FEATURE:

(D) OTHER INFORMATION: G to A transversion  
 mutation at nucleotide 316 of the D form.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ATG GGT GCA GGT GGA AGA ATG CAA GTG TCT CCT CCC TCC AAG AAG TCT	48
Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser	
1 5 10 15	
GAA ACC GAC ACC ATC AAG CGC GTA CCC TGC GAG ACA CCG CCC TTC ACT	96
Glu Thr Asp Thr Ile Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr	
20 25 30	
GTC GGA GAA CTC AAG AAA GCA ATC CCA CCG CAC TGT TTC AAA CGC TCG	144
Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser	
35 40 45	
ATC CCT CGC TCT TTC TCC TAC CTC ATC TGG GAC ATC ATC ATA GCC TCC	192
Ile Pro Arg Ser Phe Ser Tyr Leu Ile Trp Asp Ile Ile Ile Ala Ser	
50 55 60	
TGC TTC TAC TAC TAC GCC ACC ACT TAC TTC CCT CTC CTC CCT CAC CCT	240
Cys Phe Tyr Tyr Xaa Ala Thr Thr Tyr Phe Pro Leu Leu Pro His Pro	
65 70 75 80	
CTC TCC TAC TTC GCC TGG CCT CTC TAC TGG GCC TGC CAA GGG TGC GTC	288
Leu Ser Tyr Phe Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val	
85 90 95	
CTA ACC GGC GTC TGG GTC ATA GCC CAC AAG TGC GGC CAC CAC GCC TTC	336
Leu Thr Gly Val Trp Val Ile Ala His Lys Cys Gly His His Ala Phe	
100 105 110	
AGC GAC TAC CAG TGG CTT GAC GAC ACC GTC GGT CTC ATC TTC CAC TCC	384
Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser	
115 120 125	
TTC CTC CTC GTC CCT TAC TTC TCC TGG AAG TAC AGT CAT CGC AGC CAC	432
Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Ser His	
130 135 140	

2180386

CAT TCC AAC ACT GGC TCC CTC GAG AGA GAC GAA GTG TTT GTC CCC AAG His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys 145 150 155 160	480
AAG AAG TCA GAC ATC AAG TGG TAC GGC AAG TAC CTC AAC AAC CCT TTG Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu 165 170 175	528
GGA CGC ACC GTG ATG TTA ACG GTT CAG TTC ACT CTC GGC TGG CCG TTG Gly Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu 180 185 190	576
TAC TTA GCC TTC AAC GTC TCG GGA AGA CCT TAC GAC GGC GGC TTC CGT Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Arg 195 200 205	624
TGC CAT TTC CAC CCC AAC GCT CCC ATC TAC AAC GAC CGC GAG CGT CTC Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu 210 215 220	672
CAG ATA TAC ATC TCC GAC GCT GGC ATC CTC GCC GTC TGC TAC GGT CTC Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu 225 230 235 240	720
TTC CGT TAC GCC GCC GGC CAG GGA GTG GCC TCG ATG GTC TGC TTC TAC Phe Arg Tyr Ala Ala Gly Gln Gly Val Ala Ser Met Val Cys Phe Tyr 245 250 255	768
GGA GTC CCG CTT CTG ATT GTC AAT GGT TTC CTC GTG TTG ATC ACT TAC Gly Val Pro Leu Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr 260 265 270	816
TTG CAG CAC ACG CAT CCT TCC CTG CCT CAC TAC GAT TCG TCC GAG TGG Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp 275 280 285	864
GAT TGG TTC AGG GGA GCT TTG GCT ACC GTT GAC AGA GAC TAC GGA ATC Asp Trp Phe Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile 290 295 300	912
TTG AAC AAG GTC TTC CAC AAT ATT ACC GAC ACG CAC GTG GCC CAT CAT Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His 305 310 315 320	960
CCG TTC TCC ACG ATG CCG CAT TAT CAC GCG ATG GAA GCT ACC AAG GCG Pro Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala 325 330 335	1008
ATA AAG CCG ATA CTG GGA GAG TAT TAT CAG TTC GAT GGG ACG CCG GTG Ile Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Phe Asp Gly Thr Pro Val 340 345 350	1056
GTT AAG GCG ATG TGG AGG GAG GCG AAG GAG TGT ATC TAT GTG GAA CCG Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro 355 360 365	1104
GAC AGG CAA GGT GAG AAG AAA GGT GTG TTC TGG TAC AAC AAT AAG TTA T Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu 370 375 380	1153
GA	1155

- 37 -

2180386

Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp  
 275 280 285  
 Asp Trp Phe Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile  
 290 295 300  
 Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His  
 305 310 315 320  
 Pro Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala  
 325 330 335  
 Ile Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Phe Asp Gly Thr Pro Val  
 340 345 350  
 Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro  
 355 360 365  
 Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu  
 370 375 380

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1155 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Brassica napus
- (ix) FEATURE:
  - (D) OTHER INFORMATION: Wild type F form.
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

ATG GGT GCA GGT GGA AGA ATG CAA GTG TCT CCT CCC TCC AAA AAG TCT	48
Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser	
1 5 10 15	
GAA ACC GAC AAC ATC AAG CGC GTA CCC TGC GAG ACA CCG CCC TTC ACT	96
Glu Thr Asp Asn Ile Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr	
20 25 30	
GTC GGA GAA CTC AAG AAA GCA ATC CCA CCG CAC TGT TTC AAA CGC TCG	144
Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser	
35 40 45	
ATC CCT CGC TCT TTC TCC TAC CTC ATC TGG GAC ATC ATC ATA GCC TCC	192
Ile Pro Arg Ser Phe Ser Tyr Leu Ile Trp Asp Ile Ile Ile Ala Ser	
50 55 60	



2180386

TGC TTC TAC TAC GTC GCC ACC ACT TAC TTC CCT CTC CTC CCT CAC CCT Cys Phe Tyr Tyr Val Ala Thr Thr Tyr Phe Pro Leu Leu Pro His Pro 65 70 75 80	240
CTC TCC TAC TTC GCC TGG CCT CTC TAC TGG GCC TGC CAG GGC TGC GTC Leu Ser Tyr Phe Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val 85 90 95	288
CTA ACC GGC GTC TGG GTC ATA GCC CAC GAG TGC GGC CAC CAC GCC TTC Leu Thr Gly Val Trp Val Ile Ala His Glu Cys Gly His His Ala Phe 100 105 110	336
AGC GAC TAC CAG TGG CTG GAC GAC ACC GTC GGC CTC ATC TTC CAC TCC Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser 115 120 125	384
TTC CTC CTC GTC CCT TAC TTC TCC TGG AAG TAC AGT CAT CGA CGC CAC Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His 130 135 140	432
CAT TCC AAC ACT GGC TCC CTC GAG AGA GAC GAA GTG TTT GTC CCC AAG His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys 145 150 155 160	480
AAG AAG TCA GAC ATC AAG TGG TAC GGC AAG TAC CTC AAC AAC CCT TTG Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu 165 170 175	528
GGA CGC ACC GTG ATG TTA ACG GTT CAG TTC ACT CTC GGC TGG CCT TTG Gly Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu 180 185 190	576
TAC TTA GCC TTC AAC GTC TCG GGG AGA CCT TAC GAC GGC GGC TTC GCT Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Ala 195 200 205	624
TGC CAT TTC CAC CCC AAC GCT CCC ATC TAC AAC GAC CGC GAG CGT CTC Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu 210 215 220	672
CAG ATA TAC ATC TCC GAC GCT GGC ATC CTC GCC GTC TGC TAC GGT CTC Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu 225 230 235 240	720
TAC CGC TAC GCT GCT GTC CAA GGA GTT GCC TCG ATG GTC TGC TTC TAC Tyr Arg Tyr Ala Ala Val Gln Gly Val Ala Ser Met Val Cys Phe Tyr 245 250 255	768
GGA GTT CCG CTT CTG ATT GTC AAT GGG TTC TTA GTT TTG ATC ACT TAC Gly Val Pro Leu Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr 260 265 270	816
TTG CAG CAC ACG CAT CCT TCC CTG CCT CAC TAT GAC TCG TCT GAG TGG Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp 275 280 285	864
GAT TGG TTG AGG GGA GCT TTG GCC ACC GTT GAC AGA GAC TAC GGA ATC Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile 290 295 300	912

2180386

TTG AAC AAG GTC TTC CAC AAT ATC ACG GAC ACG CAC GTG GCG CAT CAC Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His 305 310 315 320	960
CTG TTC TCG ACC ATG CCG CAT TAT CAT GCG ATG GAA GCT ACG AAG GCG Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala 325 330 335	1008
ATA AAG CCG ATA CTG GGA GAG TAT TAT CAG TTG CAT GGG ACG CCG GTG Ile Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Leu His Gly Thr Pro Val 340 345 350	1056
GTT AAG GCG ATG TGG AGG GAG GCG AAG GAG TGT ATC TAT GTG GAA CCG Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro 355 360 365	1104
GAC AGG CAA GGT GAG AAG AAA GGT GTG TTC TGG TAC AAC AAT AAG TTA T Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu 370 375 380	1153
GA	1155

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 384 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser 1 5 10 15
Glu Thr Asp Asn Ile Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr 20 25 30
Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser 35 40 45
Ile Pro Arg Ser Phe Ser Tyr Leu Ile Trp Asp Ile Ile Ile Ala Ser 50 55 60
Cys Phe Tyr Tyr Val Ala Thr Thr Tyr Phe Pro Leu Leu Pro His Pro 65 70 75 80
Leu Ser Tyr Phe Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val 85 90 95
Leu Thr Gly Val Trp Val Ile Ala His Glu Cys Gly His His Ala Phe 100 105 110
Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser 115 120 125
Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His 130 135 140

2180386

His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys  
 145 150 155 160  
 Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu  
 165 170 175  
 Gly Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu  
 180 185 190  
 Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Ala  
 195 200 205  
 Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu  
 210 215 220  
 Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu  
 225 230 235 240  
 Tyr Arg Tyr Ala Ala Val Gln Gly Val Ala Ser Met Val Cys Phe Tyr  
 245 250 255  
 Gly Val Pro Leu Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr  
 260 265 270  
 Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp  
 275 280 285  
 Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile  
 290 295 300  
 Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His  
 305 310 315 320  
 Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala  
 325 330 335  
 Ile Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Leu His Gly Thr Pro Val  
 340 345 350  
 Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro  
 355 360 365  
 Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu  
 370 375 380

2180386

## WHAT IS CLAIMED IS:

1. An endogenous oil obtained from *Brassica* seeds, said oil having an oleic acid content of greater than about 80%, an  $\alpha$ -linolenic acid content of less than about 2.5% and an erucic acid content of less than about 2%, said oleic acid content, linolenic acid content and erucic acid content determined after hydrolysis of said oil.
2. The oil of claim 1, further comprising a linoleic acid content of from about 1% to about 10%, said linoleic acid content determined after hydrolysis of said oil.
3. The oil of claim 1, wherein said oleic acid content is from about 84% to about 88% and said  $\alpha$ -linolenic acid content is from about 1% to about 2%.
4. The oil of claim 3, further comprising a linoleic acid content of from about 3% to about 7%, said linoleic acid content determined after hydrolysis of said oil.
5. The oil of claim 1, wherein said seeds are *Brassica napus* seeds.
6. Crushed seeds, said seeds produced by a *Brassica* plant containing at least one recombinant nucleic construct, said at least one construct comprising:
  - a) a first seed-specific regulatory sequence fragment operably linked in sense orientation to a wild-type microsomal delta-12 fatty acid desaturase coding sequence fragment; and
  - b) a second seed-specific regulatory sequence fragment operably linked in sense orientation to a

2180386

wild-type microsomal delta-15 fatty acid desaturase coding sequence fragment,  
wherein said seeds yield an oil having an oleic acid content of about 86% or greater and an erucic acid content of less  
5 than about 2%, said oleic acid content and erucic acid content determined after hydrolysis of said oil.

7. The crushed seeds of claim 6, wherein said plant contains first and second recombinant nucleic acid constructs, said first construct comprising said delta-12  
10 desaturase coding sequence fragment and said second recombinant nucleic acid construct comprising said delta-15 desaturase coding sequence fragment.

8. The crushed seeds of claim 6, wherein said delta-12 desaturase coding sequence fragment comprises a full-length  
15 *Brassica* delta-12 desaturase coding sequence.

9. The crushed seeds of claim 6, wherein said delta-15 desaturase coding sequence fragment comprises a full-length *Brassica* delta-15 desaturase coding sequence.

10. Crushed seeds, said seeds produced by a *Brassica*  
20 plant containing at least one recombinant nucleic construct, said at least one construct comprising:

- 25 a) a first seed-specific regulatory sequence fragment operably linked to a wild-type microsomal delta-12 fatty acid desaturase coding sequence fragment; and  
b) a second seed-specific regulatory sequence fragment operably linked to a wild-type microsomal delta-15 fatty acid desaturase coding sequence fragment,

wherein said seeds yield an oil having an oleic acid content of 80% or greater, an  $\alpha$ -linolenic acid content of about 2.5% or less and an erucic acid content of less than about 2%, said oleic acid content, linolenic acid content and erucic acid content determined after hydrolysis of said oil.

11. The crushed seeds of claim 10, wherein said first and second regulatory sequence fragments are linked in sense orientation to said delta-12 and delta-15 desaturase coding sequence fragments, respectively.

10 12. The crushed seeds of claim 10, wherein said plant contains a first recombinant nucleic acid construct comprising said delta-12 desaturase coding sequence fragment and a second recombinant nucleic acid construct comprising said delta-15 desaturase coding sequence fragment.

15 13. The crushed seeds of claim 10, wherein said delta-12 desaturase coding sequence fragment comprises a full-length *Brassica* delta-12 desaturase coding sequence.

14. The crushed seeds of claim 10, wherein said delta-15 desaturase coding sequence fragment comprises a full-length *Brassica* delta-15 desaturase coding sequence.

15. The crushed seeds of claim 10, wherein said oil has an oleic acid content of about 84% to about 89%, an  $\alpha$ -linolenic acid content of about 1% to about 2% and an erucic acid content of less than about 2%, said oleic acid content, linolenic acid content and erucic acid content determined after hydrolysis of said oil.

2180386

16. The crushed seeds of claim 15, wherein said oleic acid content is from about 86% to about 89% and said  $\alpha$ -linolenic acid content is from about 1% to about 1.7%.

17. A method of producing an endogenous oil from  
5 Brassica seeds, comprising the steps of:
- a) creating at least one Brassica plant having a seed-specific reduction in microsomal delta-12 fatty acid desaturase gene expression and a seed-specific reduction in microsomal delta-15 fatty acid  
10 desaturase gene expression;
  - b) crushing seeds produced from said plant; and
  - c) extracting said oil from said seeds, said oil having an oleic acid content of about 86% or greater and an erucic acid content of less than about 2%,  
15 said oleic acid content and erucic acid content determined after hydrolysis of said oil.

18. The method of claim 17, wherein said seed-specific reduction in delta-12 desaturase expression is created by cosuppression.

- 20 19. The method of claim 17, wherein said seed-specific reduction in delta-12 desaturase expression is created by antisense suppression.

20. The method of claim 17, wherein said seed-specific reduction in delta-15 desaturase expression is created by  
25 cosuppression.

21. The method of claim 17, wherein said seed-specific reduction in delta-15 desaturase expression is created by antisense suppression.

22. A method of producing an endogenous oil from *Brassica* seeds, comprising the steps of:

5 a) creating at least one *Brassica* plant having a seed-specific reduction in microsomal delta-12 fatty acid desaturase gene expression and a seed-specific reduction in microsomal delta-15 fatty acid desaturase gene expression;

b) crushing seeds produced from said plant; and

10 c) extracting said oil from said seeds, said oil having an oleic acid content of about 80% or greater, an  $\alpha$ -linolenic acid content of 2.5% or less and an erucic acid content of less than about 2%, said oleic acid content and erucic acid content determined after hydrolysis of said oil.

15 23. The method of claim 22, wherein said seed-specific reduction in delta-12 desaturase expression is created by cosuppression.

20 24. The method of claim 22, wherein said seed-specific reduction in delta-12 desaturase expression is created by antisense suppression.

25 25. The method of claim 22, wherein said seed-specific reduction in delta-15 desaturase expression is created by cosuppression.

26. The method of claim 22, wherein said seed-specific reduction in delta-15 desaturase expression is created by antisense suppression.